

The antioxidants α -lipoic acid and *N*-acetylcysteine reverse memory impairment and brain oxidative stress in aged SAMP8 mice

Susan A. Farr,*† H. Fai Poon,‡ Dilek Dogrukol-Ak,§ Jeniffer Drake,‡ William A. Banks,* Edward Eyerman,* D. Allan Butterfield‡ and John E. Morley*

**Geriatric Research Education and Clinical Center (GRECC), VA Medical Center, St. Louis, Missouri, USA*

†*Department of Internal Medicine, Division of Geriatric Medicine, St. Louis University School of Medicine, St. Louis, Missouri, USA*

‡*Department of Chemistry, Center of Membrane Sciences, and Sander-Brown Center on Aging, University of Kentucky, Lexington, Kentucky, USA*

§*Anadolu University, Faculty of Pharmacy, Department of Analytical Chemistry, Eskisehir, Turkey*

Abstract

Oxidative stress may play a crucial role in age-related neurodegenerative disorders. Here, we examined the ability of two antioxidants, α -lipoic acid (LA) and *N*-acetylcysteine (NAC), to reverse the cognitive deficits found in the SAMP8 mouse. By 12 months of age, this strain develops elevated levels of A β and severe deficits in learning and memory. We found that 12-month-old SAMP8 mice, in comparison with 4-month-old mice, had increased levels of protein carbonyls (an index of protein oxidation), increased TBARS (an index of lipid peroxidation) and a decrease in the weakly immobilized/strongly immobilized (W/S) ratio of the protein-specific spin label MAL-6 (an index of oxidation-induced conformational changes in synaptosomal membrane proteins). Chronic administration of

either LA or NAC improved cognition of 12-month-old SAMP8 mice in both the T-maze footshock avoidance paradigm and the lever press appetitive task without inducing non-specific effects on motor activity, motivation to avoid shock, or body weight. These effects probably occurred directly within the brain, as NAC crossed the blood–brain barrier and accumulated in the brain. Furthermore, treatment of 12-month-old SAMP8 mice with LA reversed all three indexes of oxidative stress. These results support the hypothesis that oxidative stress can lead to cognitive dysfunction and provide evidence for a therapeutic role for antioxidants.

Keywords: *N*-acetylcysteine, blood–brain barrier, learning, α -lipoic acid, oxidative stress, SAMP8.
J. Neurochem. (2003) **84**, 1173–1183.

Free radical damage from oxidative stress has long been thought to play an important role in age-related neurodegenerative disorders (Harman 1995). It has been suggested that free radical damage compromises composition integrity of cell membranes, which decreases membrane fluidity (Zs-Nagy 1990). Although the specific mechanism for free radical generation and consequent oxidative stress differ between normal aging and neurodegenerative diseases, a consensus is emerging that free radical processes do play an important role in the etiology of many disorders (Hensley *et al.* 1995a; Butterfield and Stadtman 1997; Butterfield *et al.* 2001a; Butterfield and Lauderback 2002). Free radical-mediated damage to neuronal membrane components has been implicated in the etiology of diseases of aging such as Alzheimer's disease (AD). For example, the molecular basis of AD is unclear, but numerous lines of genetic and

biochemical evidence suggest that a 39–43 amino acid peptide, amyloid β peptide (A β) that is the principal component of senile plaques in the AD brain, is central to the pathogenesis of AD (Butterfield *et al.* 2001a; Butterfield *et al.* 2001b; Butterfield and Lauderback 2002). A β is

Received September 25, 2002; revised manuscript received November 22, 2002; accepted November 22, 2002.

Address correspondence and reprint requests to Dr Susan A. Farr, VA Medical Center (151/JC), 915 N. Grand Blvd., St. Louis, MO 63109, USA. E-mail: farrsa52@aol.com or Professor D. Allan Butterfield, Department of Chemistry and Center of Membrane Sciences, University of Kentucky, Lexington, KY 40506–0055, USA. E-mail: dabcsn@uky.edu

Abbreviations used: A β , amyloid β peptide; AD, Alzheimer's disease; BBB, blood–brain barrier; DNP, 2,4-dinitrophenyl hydrazone; FFA, free fatty acids; LA, α -lipoic acid; NAC, *N*-acetylcysteine; SDS, sodium dodecyl sulfate; W/S, weakly immobilized/strongly immobilized.

involved in free-radical formation that induces damage to neurons *in vitro* (Varadarajan *et al.* 2001; Yatin *et al.* 1999; Varadarajan *et al.* 2000; Kanski *et al.* 2002).

Oxidative stress can probably result in cognitive impairments. Antioxidants have been found to both prevent and reverse learning and memory deficits induced by free radicals (Bickford *et al.* 2000; Kastin *et al.* 1979; Introini *et al.* 1985; Shih *et al.* 1986; Jankovic *et al.* 1990; Guerrero *et al.* 1999; Abe and Saito 2000; Emilien *et al.* 2000; Rivas-Arancibia *et al.* 2000). α -Lipoic acid (LA) and *N*-acetylcysteine (NAC) are two antioxidants used to combat oxidative stress-induced damage (Nagamatsu *et al.* 1995; Yao *et al.* 1989; Maziere *et al.* 1999). Studies indicate that both LA and NAC protect against oxidative stress in peripheral tissues and in the central nervous system (Maziere *et al.* 1999; Yehuda and Youdim 1981; Drust and Crawford 1983; Greeley *et al.* 1989; Stoll *et al.* 1993; Packer *et al.* 1997; Martinez *et al.* 2000; Pocernich *et al.* 2002). In addition, both compounds have been found to reverse age-related impairments in memory (Martinez *et al.* 2000; Yehuda and Youdim 1981; Stoll *et al.* 1993) and LA prevents the increase in lipid peroxidase levels that occurs with age (Arivazhagan *et al.* 2000).

The SAMP8 strain of mice develops deficits in learning and memory by 12 months of age (Woods and Porte. 1983; Yagi *et al.* 1988; Flood and Morley 1998). Such deficits are not seen in other strains at this age. SAMP8 mice have elevated levels of A β caused by the overexpression of amyloid precursor protein (APP) (Morley *et al.* 2000; Cserr 1984; Kumar *et al.* 2000; Morley *et al.* 2002). Decreasing A β with antibody or antisense in 12-month-old SAMP8 mice improves learning and memory (Van Bree *et al.* 1990; Kumar *et al.* 2000; Morley *et al.* 2000). In addition, SAMP8 mice have been found to have increased free radical production in the central nervous system (Sato *et al.* 1996; Butterfield *et al.* 1997) associated with mitochondrial dysfunction (Fujibayashi *et al.* 1998). In the current study, we examined the effects of two potent antioxidants, LA and NAC, on acquisition in the T-maze footshock avoidance task and lever press appetitive task, the ability of NAC to cross the blood-brain barrier (BBB), and the ability of LA to reverse markers of oxidative stress.

Materials and methods

Subjects

Experimentally naive, 4- and 12-month-old male SAMP8 mice were obtained from our breeding colony. The colony is derived from siblings generously provided by Dr Takeda of (Kyoto University, Japan), and has been maintained as an inbred strain for 12 years under clean-room procedures (i.e. use of sterile gloves in handling mice, sterilized cages and bedding, restricted access to breeding area), and housed in microisolator HEPA filter units (Allentown Caging, PA, USA). The colony routinely undergoes serological testing for viral and bacterial contamination and has remained free of pathogens for over 5 years. Mice are housed in rooms with a

12 : 12 light/dark cycle (lights on at 06.00 h) at 20–22°C with water and food (Richmond Laboratory Rodent Diet 5001) available *ad libitum*. All experiments were conducted after institutional approval of the animal use subcommittee, which subscribes to the NIH Guide for Care and Use of Laboratory Animals.

Drug administration

LA and NAC were a gift from Jarrow Formulas, Inc. (Los Angeles, CA, USA). All drugs were dissolved in saline at pH 7.0. LA (100 mg/kg), NAC (100 mg/kg), or saline were administered subcutaneously once daily for 4 weeks at 1 PM. At the end of week one, animals were tested in the T-maze footshock avoidance paradigm. The following day, activity was tested in an open field and a 1-week habituation to milk was begun. At the end of the 1-week habituation, mice were trained in the lever press. Behavioral measures and training were conducted between 07.30 and 11.30 h.

T-maze training

Training procedures for the T-maze footshock avoidance apparatus have been described (Farr *et al.* 1999, 2000). The maze consisted of a black plastic start alley with a start box at one end and two goal boxes at the other. A stainless steel rod floor ran throughout the maze. The start box was separated from the start alley by a plastic guillotine door that prevented the mouse from moving down the alley until the training started.

A training trial began when a mouse was placed into the start box. The guillotine door was raised and the buzzer was sounded simultaneously; 5 s later, footshock was applied. The goal box the mouse first entered on the first trial was designated as 'incorrect'. Footshock was continued until the mouse entered the other goal box, which on all subsequent trials was designated 'correct' for the particular mouse. At the end of each trial, the mouse was removed from the goal box and returned to its home cage. A new trial began by placing the mouse back in the start box, sounding the buzzer, and raising the guillotine door. Footshock was applied 5 s later if the mouse did not leave the start box or failed to enter the correct goal box.

Training used an intertrial interval of 45 s and a door-bell type buzzer at 65 dB as the conditioned stimulus warning of onset of foot shock at 0.40 mA (Coulbourn Instruments scrambled grid floor shocker model E13-08). Mice were trained until they made one avoidance.

Open field

In order to eliminate the possibility that differences in acquisition were produced by changes in activity in treated mice, we examined activity in an open field. Mice were given one trial each in which they were allowed to explore freely in an open field for 15 min to determine their activity level. The open field apparatus consisted of a circular field 45 cm in diameter with sides that were 30 cm high. A testing session started with the entrance of the mouse into the side of the field facing the wall. The distances each mouse traveled during the single 15 min session were recorded in centimeters using a Polytrak recording system (San Diego Instruments).

Habituation to milk solution

Appetitive tasks used a solution of one part evaporated milk and two parts water. Mice were initially habituated to this novel food by allowing them access to it in their home cages overnight. During

three nights of habituation, food and water were removed to encourage drinking of the milk solution. After the third session, mice not consuming at least 20 mL of milk solution were excluded (less than 10% of the population). In appetitive training tasks, mice were run after overnight food and water deprivation.

Acquisition of lever press for milk reinforcement

Mice were placed into a fully automated lever press chamber. Pressing a lever on one wall of the compartment caused a light and liquid dipper with 100 μ L of milk to appear on the opposite wall. On days 1 and 2, mice had 11 s to obtain the reward; on days 3 and 4, mice had 7 s to obtain the reward; on all subsequent days, mice had 4 s to obtain a reward. Mice were given a 30-min training session each day until all groups in the particular study received an average of 100 rewards in a 30-min session. The measure of acquisition was the number of reinforced lever presses.

Blood-brain barrier studies

[14 C]N-Acetylcysteine (C-NAC) with a specific activity of 55 mCi/mmol was purchased from ICN Biomedicals, Inc. (Irvine, CA, USA). Bovine serum albumin was labeled with 99m Tc (T-Alb) by adding 0.12 mg of stannous tartrate and 1 mg of bovine serum albumin to 1-mL distilled water. The pH was adjusted to 2.5–3.3 with 0.2 M HCl. The mixture was incubated at room temperature (22°C) for 20 min before use.

Measurement of influx into brain

The unidirectional influx rate (K_i) from blood to brain was determined in mice by multiple-time regression analysis (Blasberg *et al.* 1983; Patlak *et al.* 1983; Kastin *et al.* 2001). Male ICR mice weighing 20–25 g were anesthetized with urethane and the left jugular vein and right carotid artery exposed. Mice received an injection into the jugular vein of 0.2 mL of lactated Ringer's solution with 1% bovine serum albumin and 2 (10^6) cpm of (C-NAC). Arterial blood was collected at 2, 3, 5, 7.5, 10, 15 or 20 min after intravenous injection through a cut in the carotid artery. Mice were immediately decapitated after collection of arterial blood and the whole brain removed. Whole blood was centrifuged at 5000 g at 4°C for 10 min and 50 μ L removed. The levels of radioactivity in serum and brain were determined in a beta counter. The K_i , expressed in μ L/g/min, and the apparent volume of distribution (V_i , in μ L/g) was determined from the equation

$$Am/C_{pt} = K_i \left[\int_0^t C_p(T) dT \right] / C_{pt} + V_i$$

where Am is cpm/g of brain, C_{pt} is cpm/ μ L of arterial serum at time t , and T is the dummy variable for time. Am/C_{pt} is the brain/blood ratio in μ L/g and exposure time is measured by the term $[\int_0^t C_p(T) dT] / C_{pt}$. Only the linear portion of the relation between Am/C_{pt} versus exposure time is used to compute K_i and V_i . The percentage of the intravenously injected dose taken up per gram of brain was determined from the equation:

$$\%Inj/g = 10^{-3} (Am/C_{pt}) (\%Inj/mL).$$

Capillary depletion

Mice received intravenous injections of 6.6 (10^6) cpm of (C-NAC) and 10^6 cpm of T-Alb. Ten minutes after intravenous injection, the

vascular space of the brain was washed free of blood. This washout was preceded by opening the abdomen and taking an arterial blood sample from a cut in the abdominal aorta. The jugular veins were then severed, the thorax opened, and 20 mL of lactated Ringer's solution perfused through the left ventricle of the heart while the descending thoracic aorta was occluded. Washout took less than 1 min. After washout, the mouse was immediately decapitated and the cerebral cortex removed, weighed, and homogenized with a glass homogenizer (10 strokes) in 0.8 mL of physiological buffer (10 mM Hepes, 141 mM NaCl, 4 mM KCl, 2.8 mM $CaCl_2$, 1 mM $MgSO_4$, 1 mM NaH_2PO_4 and 10 mM D-glucose adjusted to pH 7.4). Dextran solution (1.6 mL of a 26% solution) was added to the homogenate, which was vortexed and homogenized again (three strokes). Homogenization was performed on ice before centrifuging at 5400 g for 15 min at 4°C in a Beckman Allegra 21R centrifuge with a swinging bucket rotor (Fullerton, CA, USA). The pellet containing brain capillaries and the supernatant representing the brain parenchymal/brain interstitial fluid space were carefully separated and the levels of radioactivity determined in gamma and beta counters for 99m Tc and 14 C, respectively. Levels of 14 C and 99m Tc were also measured in serum. Levels of 14 C were determined after degradation of 99m Tc to non-detectable levels. The fractions were expressed as volumes of distribution (μ L/g).

Oxidative stress measures

SAMP8 brain samples were flash frozen in liquid nitrogen in St. Louis and sent to Lexington on wet ice overnight. The samples were homogenized in 0.32 M sucrose isolation buffer (2 mM EDTA, 2 mM EGTA, 20 mM Hepes, 20 μ g/mL trypsin inhibitor, 4 μ g/mL leupeptin, 4 μ g/mL pepstatin, 5 μ g/mL aprotinin) by sonication and the protein concentration was determined by the BCA method. Three indices of oxidative stress were used: (i) protein carbonyl levels, which index protein oxidation (Butterfield and Stadtman 1997); (ii) the weakly immobilized/strongly immobilized (W/S) ratio of the protein-specific spin label MAL-6, which when bound to synaptosomal membrane proteins yields a lower value of the W/S ratio, indexing oxidative stress-induced alterations in membrane protein conformation (Hensley *et al.* 1994; Butterfield and Stadtman 1997); and (iii) thiobarbituric acid reactive substance (TBARS), an index of lipid peroxidation. Protein carbonyl levels have been shown to be increased in aging (Butterfield *et al.* 1997; Hensley *et al.* 1994; Butterfield and Stadtman 1997). The W/S ratio is an EPR parameter reflective of protein-protein interactions, which decreases in oxidative stress (Hensley *et al.* 1994). The W/S ratio is the ratio of intensity of the $M_1 = +1$ low-field weakly immobilized line and $M_1 = +1$ low-field strongly immobilized resonance line of a protein-specific spin label, MAL-6 (Butterfield 1982). TBARS provide a measure of lipid peroxidation damage (Ohkawa *et al.* 1979) that was shown to be involved in aging (Tappel 1968; Harman 1969) because of the high reactivity of thiobarbituric acid with the lipid peroxidation end product, malondialdehyde (Ohkawa *et al.* 1979).

Carbonyl level

Protein carbonyl levels of proteins were determined immunochemically as adducts of 2,4-dinitrophenylhydrazine (Oliver *et al.* 1987). Five microliters of the samples were treated with an equal volume of 12% sodium dodecyl sulfate (SDS). Samples were then derivatized

with 10 μL of 20 mM 2,4-DNPH for 20 min. The reaction was stopped by addition of neutralizing reagent (7.5 μL of 2 M Tris/30% glycerol buffer, pH = 8.0). Levels of protein carbonyls were measured by using the slot-blot technique with 250 ng of protein loaded per slot. The 2,4-dinitrophenyl hydrazone (DNP) adduct of the carbonyls is detected on the nitrocellulose paper using a primary rabbit antibody (Intergen) specific for DNP-protein adducts (1 : 100) and then a secondary goat anti-rabbit IgG (Sigma) antibody. The resulting stain was developed by SCION-IMAGE software package.

TBARS

The concentration of TBARS in brain tissue was determined according to the method of Ohkawa *et al.* (1979). Fifty milliliters of 10% w/v of ice cold trichloroacetic acid was added into 0.25 mL of 4 mg/mL tissue homogenate. The samples were spun in an Eppendorf centrifuge tube for 5 min at 3000 *g*. The supernatant was collected (0.5 mL) and treated with TBA reagent (20 mM TBA in 50% v/v glacial acetic acid). The samples were then heated at 100°C for 1 h. After the cooling period, butanol was added, and the organic layer was removed and redistributed to a black microtiter plate (Corning Inc, Acton, MA, USA). End point fluorescence was measured at $\lambda_{\text{ex}} = 515 \text{ nm}$ and $\lambda_{\text{em}} = 585 \text{ nm}$.

W/S ratio

Synaptosomes were purified as described (Butterfield *et al.* 1994; Hensley *et al.* 1994; Hensley *et al.* 1995b). The crude homogenate was removed and respun at 20 000 *g* at 4°C for 10 min. The resulting pellet was resuspended in the 0.32 M sucrose isolation buffer and layered on a discontinuous sucrose gradient (10 mL of 1.18 M sucrose, pH 8.5/10 mL of 1.0 M sucrose, pH 7.4/10 mL of 0.85 M sucrose, pH 7.4, each containing 2 mM EDTA/2 mM EGTA/10 mM Hepes). The samples were then spun at 82 500 *g* at 4°C for 60 min in a Beckman swinging-bucket rotor. Synaptosomes were removed from the 1.18/1.0 M sucrose interface and resuspended in 20 mL of lysing buffer (10 mM Hepes/2 mM EDTA/2 mM EGTA, pH 7.4). The samples were then centrifuged at 32 000 *g* at 4°C for 10 min. The pellet was removed and resuspended in PBS buffer and spun down twice more. After the third wash, the protein concentration was determined by the BCA method.

Spin labeling of synaptosomal membrane proteins was performed as described (Umhauer *et al.* 1992; Hensley *et al.* 1994). Isolated synaptosomes were suspended in lysing buffer for 30 min. Lysed synaptosomal membranes were labeled with the protein-specific spin label MAL-6. After incubation for 18 h at 4°C with 20 μg MAL-6/mg protein, samples were washed six times in lysing buffer to remove excess spin label. The pellet was then resuspended in approximately 400 μL lysing buffer and allowed to come to room temperature. EPR spectra were acquired on a Bruker model EMX EPR spectrometer (Bruker, Billerica, MA, USA) operating at an incident microwave power of 18 mW, a modulation amplitude of 0.4 G, a time constant of 1.28 ms, and a conversion time of 10 ms.

Statistics

For behavioral studies, results are expressed as means with their standard errors. All groups had 10 mice. The acquisition test scores for T-maze (mean trials to make first avoidance) and lever press (number of reinforced lever presses) for each group were analyzed

with one-way analysis of variance (ANOVA) and a two-way analysis of variance, respectively. Latencies to escape shock, activity, food intake, and weight change were analyzed by a one-way ANOVA. The control and treatment groups were compared with Tukey's *t*-test or Tukey's HSD (Keppel and Zedeck 1989).

For BBB pharmacokinetic analysis, regression lines were calculated by the least squares method and compared statistically with the PRISM 3.0 program (GraphPad Software, Inc, San Diego, CA, USA). Regression lines are reported with their slope, standard error of the mean, correlation coefficient (*r*), the number of mice per line (*n*), and the level of the statistical significance (*p*). Means are reported with their standard error and *n*.

Oxidative stress measures were analyzed by Student's *t*-tests. A value of *p* < 0.05 was considered statistically significant.

Results

Effects of LA on cognition

Administration of LA improved acquisition as tested in the T-maze footshock avoidance paradigm (Fig. 1a). The ANOVA for the trials to first avoidance measure indicated a significant effect ($F_{2,27} = 34.31$, *p* < 0.001). Tukey's *t*-test *post-hoc* analysis revealed 12-month-old SAMP8 mice that had received LA took significantly fewer trials to reach criterion than the 12-month-old SAMP8 mice which received saline. The group of mice administered LA did not significantly differ from the 4-month-old SAMP8 mice administered saline. The ANOVA for latencies to escape shock on the first trial did not indicate a significant difference.

The ANOVA for activity in an open field indicated a statistically significant effect ($F_{2,27} = 11.48$, *p* < 0.001). Tukey's HSD *post-hoc* analysis indicated that the 4-month-old SAMP8 mice ($1897 \pm 65 \text{ cm}$, *n* = 10) were significantly more active in an open field than the 12-month-old SAMP8 mice which received either saline ($1583 \pm 42 \text{ cm}$, *n* = 10) or LA ($1656 \pm 33 \text{ cm}$, *n* = 11). The groups of 12-month-old SAMP8 mice receiving either saline or LA did not differ from one another.

The two-way ANOVA for the number of rewarded lever presses showed a significant effect for group ($F_{2,260} = 50.86$, *p* < 0.0001), day ($F_{9,260} = 42.80$, *p* < 0.0001), and the interaction group \times day ($F_{18,260} = 4.18$, *p* < 0.0001). Tukey's *post-hoc* analysis indicated that 12-month-old SAMP8 mice administered LA achieved significantly more rewards on days 7–9 compared with the mice administered saline (Fig. 1b). Four-month-old SAMP8 mice achieved significantly more rewards than either of the 12 month groups on days 7–10.

The ANOVA analyzing food intake in the mice administered LA indicated a statistically significant effect ($F_{2,27} = 3.62$, *p* < 0.05). Tukey's HSD *post-hoc* analysis indicated that the 12-month-old SAMP8 mice administered LA ate significantly more than the 4-month-old SAMP8 mice.




Fig. 1 Effects of LA and NAC on acquisition of T-maze footshock avoidance and lever press. (a) LA improved acquisition of T-maze footshock avoidance in 12-month-old SAMP8 mice. *Indicates that the value differs from the saline treated 12-month-old SAMP8 mice at $p < 0.01$. (b) LA improved acquisition of lever press in 12-month-old SAMP8 mice. On days 7–9 of training, 12-month-old SAMP8 mice receiving LA achieved more rewards than the day-matched 12-month-old SAMP8 mice which received saline $p < 0.01$. Four-month-old SAMP8 mice achieved more rewards than either 12-month-old SAMP8 groups on days 7–10 $p < 0.01$. * $p < 0.01$ in comparison to

day-matched 12-month-old SAMP8 mice receiving saline. (c) NAC improved acquisition of T-maze footshock avoidance in 12-month-old SAMP8 mice. *Indicates that the value differs from the saline treated 12-month-old SAMP8 mice at $p < 0.01$. (d) NAC improved acquisition of lever press in 12 month SAMP8 mice. On day 7, 12-month-old SAMP8 mice receiving NAC achieved more rewards than the 12-month-old SAMP8 mice receiving saline * $p < 0.01$. Four-month-old SAMP8 mice achieved more rewards than both 12-month-old SAMP8 groups on days 7–9, $p < 0.01$.

Twelve-month-old SAMP8 mice administered saline were not significantly different from the 12-month-old SAMP8 mice administered LA nor the 4-month-old SAMP8 mice. The ANOVA for body weight change during treatment did not show a significant effect (see Table 1).

Effects of NAC on cognition

The ANOVA analyzing T-maze acquisition showed a significant effect ($F_{2,25} = 61.29, p < 0.001$). Tukey's *post-hoc* analysis showed that the 12-month-old SAMP8 mice which

received NAC performed significantly better than the 12-month-old SAMP8 mice which received saline and were not different from the 4 month SAMP8 mice (Fig. 1c). The ANOVA for latencies to escape shock on first trial did not show a significant effect.

The ANOVA for activity showed a significant effect ($F_{2,26} = 4.07, p < 0.05$). Tukey's *post-hoc* analysis showed the only difference to be that the 4-month-old SAMP8 mice administered saline ($1897 \pm 65, n = 10$) were significantly more active than the 12-month-old SAMP8 mice

Table 1 Effects of antioxidants on food intake and body weight

	Average daily food intake (g)	Weight change (g)
Study 1		
12 M P8 saline	5.11 ± 0.83 ^{a,b}	+ 0.07 ± 0.90 ^a
12 M P8 α -lipoic	5.78 ± 0.76 ^a	- 0.36 ± 1.22 ^a
4 M P8 saline	4.93 ± 0.63 ^b	- 0.54 ± 0.62 ^a
Study 2		
12 M P8 saline	5.79 ± 0.84 ^{a,b}	+ 0.40 ± 0.47 ^{a,b}
12 M P8 <i>N</i> -acetylcysteine	6.07 ± 0.71 ^a	+ 0.88 ± 0.72 ^a
4 M P8 saline	4.99 ± 0.55 ^b	- 0.46 ± 0.84 ^b

M refers to age in months; P8 refers to SAMP8 mice. ^{a,b}Groups with different letters are significantly differ $p < 0.05$ for that study.

administered NAC (1668 ± 66 cm, $n = 10$). Twelve-month-old-mice given saline (1720 ± 52 cm, $n = 9$) were not different from 12-month-old mice given NAC.

The two-way ANOVA for rewarded presses in the lever press study showed a significant effect for group ($F_{2,234} = 28.76$, $p < 0.0001$), day ($F_{8,234} = 47.00$, $p < 0.0001$), and the interaction group × day ($F_{2,8} = 2.42$, $p < 0.002$). Tukey's *post-hoc* analysis indicated that 12-month-old SAMP8 mice administered NAC achieved significantly more rewards on

day 7 compared with the 12-month-old SAMP8 mice administered saline (Fig. 1d). Four-month-old SAMP8 mice achieved significantly more rewards than both 12-month-old groups on the other days.

The ANOVA for food intake did not show a significant effect. The ANOVA for body weight change showed a statistically significant difference ($F_{2,27} = 9.57$, $p < 0.001$). Tukey's HSD *post-hoc* analysis indicated that the 12-month-old SAMP8 mice which received saline and NAC gained weight, whereas the 4-month-old SAMP8 mice lost weight (see Table 1).

Blood-brain barrier permeability to NAC

Figure 2(a) shows the relation between the log of levels of radioactivity in arterial serum expressed as the percentage/milliliter of injected dose (%Inj/mL) versus time after the intravenous injection of C-NAC. This relation had a slope of -0.0222 and an intercept 1.051 ($n = 13$, $r = 0.908$, $p < 0.0001$). This gave a half-time disappearance from blood of 13.5 min and a volume of distribution of 8.89 mL.

Figure 2(b) shows the relation between the brain/blood ratios (Am/Cpt) and exposure time for mice which received C-NAC. This relation was statistically significant ($r = 0.955$, $n = 13$, $p < 0.0001$) with $K_i = 2.41 \pm 0.226$ $\mu\text{L/g-min}$ and $V_i = 39.3 \pm 2.79$ $\mu\text{L/g}$. The percentage of the injected dose




Fig. 2 (a) Clearance of NAC from blood after intravenous injection. Half-time disappearance from blood was 13.5 min and the volume of distribution was 8.89 mL. (b) Multiple-time regression analysis of NAC transport across the BBB. The unidirectional influx rate was measured to be 2.41 ± 0.226 $\mu\text{L/g-min}$. (c) The percentage of an intravenously

injected dose taken up per gram of brain (%Inj/g) for NAC. Values were about 0.4 %Inj/g. (d) Capillary depletion. Results show that most of the NAC taken up by brain completely crossed the BBB to enter the parenchymal space of the brain.

taken up per gram of brain is shown in Fig. 2(c) as determined from the equation:

$$\%Inj/g = 10^{-3} (Am/Cpt)(\%Inj/mL).$$

These results show that about 0.4 %Inj/g was taken up by brain.

C-NAC crossed the endothelial barrier of the cerebral cortex as shown by recovery of radioactivity from the parenchymal space of the brain (Fig. 2d). About 2.5 μ L/g of C-NAC was present in capillaries and may reflect uptake by or binding to endothelium. For parenchyma, about 12.5 μ L of C-NAC was present, indicating that the majority of C-NAC crossed the BBB.

Measures of oxidative stress

Protein carbonyl levels

To determine whether the protein carbonyl levels of cortical synaptosomal membranes were increased in the 12-month-old SAMP8 mice relative to those from 4-month-old SAMP8 mice, the 2,4-dinitrophenylhydrazine adducts were measured immunochemically. The results showed that the protein carbonyl levels of brain proteins in 12-month-old SAMP8 were significantly greater than those of 4-month-old SAMP8 by 29% (Fig. 3aA; $p < 0.01$). However, treatment with LA reduced this difference to a statistically non-significant 17% increase (Fig. 3aB).

W/S ratio

Consistent with the result for protein carbonyl levels, the W/S ratio was decreased significantly in the synaptosomal membrane proteins from 12-month-old SAMP8 mice when compared with 4-month-old SAMP8 mice (Fig. 3bA; $p < 0.001$). As noted above, the W/S ratio is lowered by oxidative stress (Hensley *et al.* 1994; Hall *et al.* 1995a; Hall *et al.* 1995b; Hall *et al.* 1995c; Butterfield *et al.* 1997). Treatment with LA abolished the difference between aged and young SAMP8 mice (Fig. 3bB).

TBARS

Increased TBARS levels were observed in the brains of 12-month-old SAMP8 mice when compared with 4-month-old SAMP8 mice (Fig. 3cA; $p < 0.05$). This result suggested an increased lipid peroxidation in 12-month-old SAMP8 mice. Treatment with LA abolished the difference between young and old mice (Fig. 3cB).

Discussion

Oxidative stress and damage induced by free radicals has been proposed as an important mechanism for both normal aging and the cognitive decline of neurodegenerative diseases (Zs-Nagy 1978; Zs-Nagy 1990). Here, we investigated




Fig. 3 (a) Protein carbonyl levels for 4 and 12-month-old SAMP8 mice. A, Untreated mice. Error bars indicate SEM for eight animals in each group. B, LA treated mice. Error bars indicate SEM for seven animals in each group. Measured value is normalized with the 4-month-old values. * $p < 0.01$. (b) W/S ratios for 4 and 12-month-old SAMP8 mice. A, Untreated mice. Error bars indicate SEM for six and seven animals, respectively. B, LA-treated mice. Error bar indicates SEM for six animals in each group. Measured value is normalized to the 4-month-old values. * $p < 0.001$. (c) TBARS of 4 and 12-month-old SAMP8 mice. (i) Untreated mice. Error bars indicate SEM for seven and six animals, respectively. (ii) LA treated mice. Error bar indicates SEM for seven and six animals in each group. Measured value is normalized with the 4-month-old values. * $p < 0.01$.

whether antioxidant treatment could reverse the cognitive decline and oxidative damage seen in the aged SAMP8 mouse. This strain of mouse starts at about 6–8 months of age to overexpress A β , the peptide postulated to be the cause of AD (Kumar *et al.* 2000; Morley *et al.* 2000). With A β overexpression, these mice develop cognitive impairments which precedes by several months physical decline (Takeda

et al. 1991; Flood and Morley 1993; Nomura *et al.* 1996; Tanaka *et al.* 1998; Butterfield *et al.* 2001b; Morley *et al.* 2002). By 12 months of age, SAMP8 mice are healthy but have developed severe impairments in learning and memory, which are reversed by antibody or antisense directed at A β (Kumar *et al.* 2000; Morley *et al.* 2000; Banks *et al.* 2001). Other strains of mice at this age show little or no cognitive decline.

Here, we found that both LA and NAC could reverse impaired learning in 12-month-old SAMP8 mice in two separate behavioral paradigms. In the footshock avoidance T-maze, LA and NAC each returned learning to a level that was not different from unimpaired 4-month-old SAMP8 mice. LA and NAC were less effective in the lever press test. Although either antioxidant improved performance in this food-reward paradigm to a statistically significant degree, 12-month-old SAMP8 mice still underperformed when compared with 4-month-old mice.

The reason why antioxidants were not as potent in a food-reward-based learning paradigm is not clear, but one possibility is that anorexia of aging, which is thought to be unrelated to free radical production (Morley 2001; Morley 1997). Antioxidant treatment did not affect motor activity, response to shock, or body weight. Effects on these parameters could have acted as confounders and an absence of effect further supports a direct action of antioxidants on cognition.

These results are similar to those that found diets rich in antioxidants improved learning in aged rats (Bickford *et al.* 2000). In addition, two of three diets tested significantly increased brain concentrations of glutathione, the most potent known intracellular antioxidant. LA and NAC are precursors to glutathione (Overton and Fisher 1991; Wernerman and Hammarqvist 1996; Exner *et al.* 2000). The cognitive impairments of the SAMP8 mouse, however, are much greater than those occurring with normal aging.

The ability of antioxidants to reverse cognitive impairments depends upon their ability to reach the brain. This, in turn, depends on an ability to cross the BBB. The accumulation of free fatty acids (FFA) in brain results from a complex interplay between free and serum protein bound FFA, BBB influx transporters, BBB efflux transporters, and brain utilization (Banks *et al.* 1997b; Rapoport and Robinson 1995). Amino acid uptake by the brain depends on BBB transporters such as the one for large neutral amino acids (Davson and Segal 1996). Here, we showed that NAC entered the brain at a rate of about 2.41 $\mu\text{L/g}\cdot\text{min}$, a modest rate in comparison to essential amino acids and about the same rate as many centrally active peptides (Begley 1994; Banks *et al.* 1995a; Banks *et al.* 1995b; Banks *et al.* 1997a; Banks 1999). Results of the capillary depletion experiment showed NAC could completely cross the capillary wall to enter the brain tissue and extracellular space. To determine the percentage of an intravenous dose of NAC taken up by a g of brain (%*Inj/g*),

we first had to calculate circulating pharmacokinetic parameters (shown in Fig. 5). These values and the influx characteristics were then used to calculate %*Inj/g*. The results showed that about 0.4% of an injected dose was taken up by brain. This is about 4, 5, 20, and 200 times greater than the values for acetaminophen, interleukin-1 alpha, morphine, and domoic acid, respectively (Banks *et al.* 1991; Preston and Hynie 1991; Banks and Kastin 1994; Courade *et al.* 2001), all CNS-active agents. Therefore, the amount of NAC transported across the BBB is well within the therapeutic range of compounds known to exert effects on the brain.

We assessed the effect of antioxidant treatment on measures of oxidative stress in the aged SAMP8 mouse. We found that 12-month-old SAMP8 mice had more oxidative stress than 4-month-old SAMP8 mice. This confirms previous findings from another colony of SAMP8 mice and is consistent with studies showing that the SAMP8 mouse accumulates more oxidative damage than animals which age normally (Butterfield *et al.* 1997; Stadtman 1992; Hensley *et al.* 1994; Fujibayashi *et al.* 1998). Additionally, the brains of aged SAMP8 mice have a 44–50% decrease in delta-9 desaturase activity and a corresponding decrease in unsaturated free fatty acids. (Butterfield *et al.* 2001a).

The results suggest that oxidative stress is widespread, affecting measures of protein oxidation, lipid peroxidation, and oxidation-dependent changes in membrane protein conformation. These parameters were all reversed by treatment with LA. Ames and coworkers have shown that LA is able to partially reverse memory loss in normal aging rats by delaying mitochondrial dysfunction and RNA/DNA oxidation (Liu *et al.* 2002). Mitochondrial dysfunction is accompanied by a leakage into cytoplasm of O₂ and H₂O₂. That LA is readily taken up into mitochondria where it acts as a cofactor in oxidative decarboxylation of α -keto acids has led to the view that LA maybe a useful therapeutic agent in diseases characterized by mitochondria dysfunction or oxidative stress (Packer *et al.* 1997; Lynch 2001).

LA derives its antioxidant capability from its ability to (i) act as a scavenger of reactive oxygen species (ROS); (ii) chelate metals; and (iii) recycle endogenous antioxidants (Lynch 2001). LA can scavenge singlet oxygen, H₂O₂, OH \cdot , NO and ONOO $^-$. The reduced form of LA, dihyrolipoic acid, can further scavenge O₂ and peroxy radicals (Kagan *et al.* 1992). LA can also chelate several divalent cations, e.g. Mn²⁺, Cu²⁺, Zn²⁺, Cd²⁺, Pb²⁺. Therefore, LA can inhibit ascorbate-induced production of H₂O₂ by Cu⁺ (Ou *et al.* 1995). LA can recycle endogenous antioxidants, such as GSH (Ou *et al.* 1995) and vitamin C (Drake *et al.* 2002), which regenerate vitamin E. GSH, vitamin C, and vitamin E all protect the brain from oxidative stress (Drake *et al.* 2002).

In conclusion, we found treatment with the antioxidants LA and NAC reversed the age-related cognitive impairment in SAMP8 mice. These substances probably act directly on

the brain, as NAC crossed the BBB and accumulated in brain to a significant degree. Treatment with LA reversed the oxidative stress seen in 12-month-old SAMP8 mice to levels that were not different from those seen in 4-month-old SAMP8 mice. These results support the hypothesis that oxidative stress can lead to cognitive dysfunction and provide evidence for a therapeutic role for antioxidants.

Acknowledgements

This research was supported by the Medical Research Service of the Department of Veterans Affairs (Merit Review), R01 NS41863, and R01 AA12743.

References

- Abe K. and Saito H. (2000) Effects of saffron extract and its constituent crocin on learning behavior and long-term potentiation. *Phytotherapy Res.* **14**, 149–152.
- Arivazhagan P., Juliet P. and Panneerselvam C. (2000) Effect of dl-alpha-lipoic acid peroxidation and antioxidants in aged rats. *Pharm. Res.* **41**, 299–303.
- Banks W. A. (1999) Physiology and pathophysiology of the blood–brain barrier: implications for microbial pathogenesis, drug delivery and neurodegenerative disorders. *J. Neurovirol.* **5**, 538–555.
- Banks W. A. and Kastin A. J. (1994) Opposite direction of transport across the blood–brain barrier for Tyr-MIF-1 and MIF-1: comparison with morphine. *Peptides* **15**, 23–29.
- Banks W. A., Ortiz L., Plotkin S. R. and Kastin A. J. (1991) Human interleukin (IL) 1 α , murine IL-1 α and murine IL-1 β are transported from blood to brain in the mouse by a shared saturable mechanism. *J. Pharmacol. Exp Ther* **259**, 988–996.
- Banks W. A., Kastin A. J. and Jaspan J. B. (1995a) Regional variation in transport of pancreatic polypeptide across the blood–brain barrier of mice. *Pharmacol. Biochem. Behav.* **51**, 139–147.
- Banks W. A., Wustrow D. J., Cody W. L., Davis M. D. and Kastin A. J. (1995b) Permeability of the blood–brain barrier to the neurotensin_{8–13} analog NT1. *Brain Res.* **695**, 59–63.
- Banks W. A., Jaspan J. B., Huang W. and Kastin A. J. (1997a) Transport of insulin across the blood–brain barrier: saturability at euglycemic doses of insulin. *Peptides* **18**, 1423–1429.
- Banks W. A., Kastin A. J. and Rapoport S. I. (1997b) Permeability of the blood–brain barrier to circulating free fatty acids. In: *Handbook of Essential Fatty Acid Biology: Biochemistry, Physiology, and Behavioral Neurobiology* (Yehuda S. and Mostofsky D. I., eds), pp. 3–14. Human Press, Totowa, NJ.
- Banks W. A., Farr S. A., Butt W., Kumar V. B., Franko M. W. and Morley J. E. (2001) Delivery across the blood–brain barrier of antisense directed against amyloid β : reversal of learning and memory deficits in mice overexpressing amyloid precursor protein. *J. Pharmacol. Exp Ther* **297**, 1113–1121.
- Begley D. J. (1994) Strategies for delivery of peptide drugs to the central nervous system: exploiting molecular structure. *J. Control. Release* **29**, 293–306.
- Bickford P. C., Gould T., Briederick L., Chadman K., Pollock A., Young D., Shukitt-Hale B. and Joseph J. (2000) Antioxidant-rich diets improve cerebellar physiology and motor learning in aged rats. *Brain Res.* **866**, 211–217.
- Blasberg R. G., Fenstermacher J. D. and Patlak C. S. (1983) Transport of α -aminoisobutyric acid across brain capillary and cellular membranes. *J. Cereb. Blood Flow Metab.* **3**, 8–32.
- Butterfield D. A. (1982) Spin labeling in disease. In: *Biological Magnetic Resonance*, Vol. IV (Berliner L. J. and Reuben J., eds), pp. 1–78. Plenum Press, New York.
- Butterfield D. A. and Lauderback C. M. (2002) Lipid peroxidation and protein oxidation in Alzheimer's disease brain: potential causes and consequences involving amyloid beta-peptide-associated free radical oxidative stress. *Free Rad. Biol. Med.* **32**, 1050–1060.
- Butterfield D. A. and Stadtman E. R. (1997) Protein oxidation processes in aging brain. *Adv. Cell Aging Gerontol.* **2**, 161–191.
- Butterfield D. A., Hensley Hensley K., Harris M., Mattson M. and Carney J. (1994) beta-Amyloid peptide free radical fragments initiate synaptosomal lipoperoxidation in a sequence-specific fashion: implications to Alzheimer's disease. *Biochem. Biophys. Res. Commun.* **200**, 710–715.
- Butterfield D. A., Howard B. J., Yatin S., Allen K. L. and Carney J. M. (1997) Free radical oxidation of brain proteins in accelerated senescence and its modulation by *N-tert-butyl- α -phenylnitron*. *Proc. Natl Acad. Sci. USA* **94**, 674–678.
- Butterfield D. A., Drake J., Pocernich C. and Castegna A. (2001a) Evidence of oxidative damage in Alzheimer's disease brain: central role for amyloid beta-peptide. *Trends Mol Med.* **7**, 548–554.
- Butterfield D. A., Howard B. J. and LaFontaine M. A. (2001b) Brain oxidative stress in animal models of accelerated aging and the age-related neurodegenerative disorders, Alzheimer's disease and Huntington's disease. *Curr. Med. Chem.* **8**, 815–828.
- Courade J. P., Besse d'Delchambre C., Hanoun N., Hamon M., Eschaliere A., Caussade F. and Cloarec A. (2001) Acetaminophen distribution in the rat central nervous system. *Life Sci.* **69**, 1455–1464.
- Cserr H. F. (1984) Convection of brain interstitial fluid. In: *Hydrocephalus* (Shapiro K., Marmarou A. and Portnoy H., eds), pp. 59–68. Raven Press, New York.
- Davson H. and Segal M. B. (1996) Special aspects of the blood–brain barrier, in *Physiology of the CSF and Blood–Brain Barriers*, pp. 303–485. CRC Press, Boca Raton.
- Drake J., Kanski J., Varadarajan S., Tsoras M. and Butterfield D. A. (2002) Elevation of brain glutathione by gamma-glutamylcysteine ethyl ester protects against peroxynitrite-induced oxidative stress. *J. Neurosci. Res.* **68**, 776–784.
- Drust E. G. and Crawford I. L. (1983) Comparison of the effects of TRH and D-Ala²-met-enkephalinamide on hippocampal electrical activity and behavior in the unanesthetized rat. *Peptides* **4**, 239–243.
- Emilien G., Beyreuther K., Masters C. L. and Maloteaux J. M. (2000) Prospects for pharmacological intervention in Alzheimer disease. *Arch. Neurol.* **57**, 459.
- Exner R., Wessner B., Manhart N. and Roth E. (2000) Therapeutic potential of glutathione. *Wien Klin. Wochensh.* **112**, 610–616.
- Farr S. A., Uezu K., Flood J. F. and Morley J. E. (1999) Septo-hippocampal drug interactions in post-trial memory processing. *Brain Res.* **847**, 221–230.
- Farr S. A., Banks W. A., La Scola M. E., Flood J. F. and Morley J. E. (2000) Permanent and temporary inactivation of the hippocampus impairs T-maze footshock avoidance acquisition and retention. *Brain Res.* **872**, 242–249.
- Flood J. F. and Morley J. E. (1993) Age-related changes in footshock avoidance acquisition and retention in senescence accelerated mouse (SAM). *Neurobiol. Aging* **14**, 153–157.
- Flood J. F. and Morley J. E. (1998) Learning and memory in the SAMP8 mouse. *Neurosci. Biobehav. Rev.* **22**, 1–20.
- Fujibayashi Y., Yamamoto S., Waki A., Konishi J. and Yonekura Y. (1998) Increased mitochondrial DNA deletion in the brain of SAMP8, a mouse model for spontaneous oxidative stress brain. *Neurosci. Lett.* **254**, 109–112.

- Greeley G. H., Cooper C. W., Jeng Y. J., Eldridge C. and Thompson J. C. (1989) Intracerebroventricular administration of calcitonin enhances glucose-stimulated release of insulin. *Reg. Pept.* **24**, 259–268.
- Guerrero A. L., Dorado-Martinez C., Rodriguez A., Pedroza-Rios K., Borgonio-Perez G. and Rivas-Arancibia S. (1999) Effects of vitamin E on ozone-induced memory deficits and lipid peroxidation in rats. *Neuroreport* **10**, 1692.
- Hall N. C., Carney J. M., Cheng M. and Butterfield D. A. (1995a) Prevention of ischemia/reperfusion-induced alterations in synaptosomal membrane-associated proteins and lipids by *N*-tert-butyl-alpha-phenylnitron and difluoromethylornithine. *Neuroscience* **69**, 591–600.
- Hall N. C., Carney J. M., Cheng M. S. and Butterfield D. A. (1995b) Ischemia/reperfusion-induced changes in membrane proteins and lipids of gerbil cortical synaptosomes. *Neuroscience* **64**, 81–89.
- Hall N. C., Dempsey R. J., Carney J. M., Donaldson D. L. and Butterfield D. A. (1995c) Structural alterations in synaptosomal membrane-associated proteins and lipids by transient middle cerebral artery occlusion in the cat. *Neurochem. Res.* **20**, 1161–1169.
- Harman D. (1969) Prolongation of life: role of free radical reactions in aging. *J. Am. Geriatr. Soc.* **17**, 721–735.
- Harman D. (1995) Free radical theory of aging: Alzheimer's disease pathogenesis. *Age* **18**, 97–119.
- Hensley K., Carney J., Hall N., Shaw W. and Butterfield D. A. (1994) Electron paramagnetic resonance investigations of free radical-induced alterations in neocortical synaptosomal membrane protein infrastructure. *Free Rad. Biol. Med.* **17**, 321–331.
- Hensley K., Hall N., Sburamian R., Cole P., Harris M., Aksenov M., Aksenova M., Gabbita S. P., Wu J. F., Carney J. M., Lovell M., Markesbery W. R. and Butterfield D. A. (1995a) Brain regional correspondence between Alzheimer's disease histopathology and biomarkers of protein oxidation. *J. Neurochem.* **65**, 2146–2156.
- Hensley K., Howard B. J., Carney J. M. and Butterfield D. A. (1995b) Membrane protein alterations in rodent erythrocytes and synaptosomes due to aging and hyperoxia. *Biochim. Biophys. Acta* **1270**, 203–206.
- Introni I. B., McGaugh J. L. and Baratti C. M. (1985) Pharmacological evidence of a central effect of naltrexone, morphine, and β -endorphin and a peripheral effect of met- and leu-enkephalin on retention of an inhibitory response in mice. *Behav. Neural Biol.* **44**, 434–446.
- Jankovic B. D., Maric D. and Veljic J. (1990) Cerebrally mediated modulation of anaphylactic shock by methionine-enkephalin. *Int. J. Neurosci.* **51**, 193–194.
- Kagan V. E., Shvedova A., Serbinova E., Khan S., Swanson C., Powell R. and Packer L. (1992) Dihydrolipoic acid – a universal antioxidant both in the membrane and in the aqueous phase. Reduction of peroxy, ascorbyl and chromanoxyl radicals. *Biochem. Pharmacol.* **44**, 1637–1649.
- Kanski J., Aksenova M., Schoneich C. and Butterfield D. A. (2002) Substitution of isoleucine-31 by helical-breaking proline abolishes oxidative stress and neurotoxic properties of Alzheimer's amyloid beta-peptide. *Free Rad. Biol. Med.* **32**, 1205–1211.
- Kastin A. J., Olson R. D., Schally A. V. and Coy D. H. (1979) CNS effects of peripherally administered brain peptides. *Life Sci.* **25**, 401–414.
- Kastin A. J., Akerstrom V. and Pan W. (2001) Validity of multiple-time regression analysis in measurement of tritiated and iodinated leptin crossing the blood–brain barrier: meaningful controls. *Peptides* **22**, 2127–2136.
- Keppel G. and Zedeck S. (1989) *Data Analysis for Research Designs: Analysis of Variance and Multiple Regression/Correlation Approaches*. W.H. Freeman, New York.
- Kumar V. B., Farr S. A., Flood J. F., Kamlesh V., Franko M., Banks W. A. and Morley J. E. (2000) Site-directed antisense oligonucleotide decreases the expression of amyloid precursor protein and reverses deficits in learning and memory in aged SAMP8 mice. *Peptides* **21**, 1769–1775.
- Liu J., Head E., Gharib A. M., Yuan W., Ingersoll R. T., Hagen T. M., Cotman C. W. and Ames B. N. (2002) Memory loss in old rats is associated with brain mitochondrial decay and RNA/DNA oxidation: partial reversal by feeding acetyl-L-carnitine and/or R-alpha-lipoic acid. *Proc. Natl Acad. Sci. USA* **99**, 2356–2361.
- Lynch M. A. (2001) Lipoic acid confers protection against oxidative injury in non-neuronal and neuronal tissue. *Nutrit. Neurosci.* **4**, 419–438.
- Martinez M., Hernandez A. I. and Martinez N. (2000) *N*-Acetylcysteine delays age-associated memory impairment in mice. *Brain Res.* **855**, 100–106.
- Maziere C., Conte M., Degonville J., Ali D. and Maziere J. (1999) Cellular enrichment with polyunsaturated fatty acids induces an oxidative stress and activates the transcription factors AP1 and NF κ B. *Biochem. Biophys. Res. Commun.* **265**, 116–122.
- Morley J. E. (1997) Anorexia of aging: physiologic and pathologic. *Am. J. Clin. Nutr.* **66**, 760–773.
- Morley J. E. (2001) Decreased food intake with aging. *J. Gerontol. [A]* **56A**, 81–88.
- Morley J. E., Kumar V. B., Bernardo A. F., Farr S. A., Uezu K., Tumosa N. and Flood J. F. (2000) β -Amyloid precursor polypeptide in SAMP8 mice affects learning and memory. *Peptides* **21**, 1761–1767.
- Morley J. E., Farr S. A., Kumar V. B. and Banks W. A. (2002) Alzheimer's disease through the eye of a mouse: acceptance lecture for the 2001 Gayle A. Olson and Richard D. Olson prize. *Peptides* **23**, 589–599.
- Nagamatsu M., Nickander K. K., Schmelzer J. D., Raya A., Wittrock D. A., Tritschler H. and Low P. A. (1995) Lipoic acid improves nerve blood flow, reduces oxidative stress, and improves distal nerve conduction in experimental diabetic neuropathy. *Diabetes Care* **18**, 1160–1167.
- Nomura Y., Yamanaka Y., Kitamura Y., Arima T., Ohnuki T., Oomura Y., Sasaki K., Nagashima K. and Ihara Y. (1996) Senescence-accelerated mouse. Neurochemical studies on aging. *Ann. NY Acad. Sci.* **786**, 410–418.
- Ohkawa H., Ohishi N. and Yagi K. (1979) Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. *Anal. Biochem.* **95**, 351–358.
- Okuda C., Tanaka H. and Miyazaki M. (1988) Cardiovascular effect of intravenously administered thyrotropin-releasing hormone and its concentration in push-pull perfusion of the fourth ventricle in conscious and pentobarbital-anesthetized rats. *Life Sci.* **42**, 1181–1188.
- Oliver C. N., Ahn B. W., Moerman E. J., Goldstein S. and Stadtman E. R. (1987) Age-related changes in oxidized proteins. *J. Biol. Chem.* **262**, 5488–5491.
- Ou P., Tritschler H. J. and Wolff S. P. (1995) Thioctic (lipoic) acid: a therapeutic metal-chelating antioxidant? *Biochem. Pharmacol.* **50**, 123–126.
- Overton J. M. and Fisher L. A. (1991) Differentiated hemodynamic responses to central versus peripheral administration of corticotropin-releasing factor in conscious rats. *J. Autonomic Nerv. Syst.* **35**, 43–52.
- Packer L., Tritschler H. J. and Wessel K. (1997) Neuroprotection by the metabolic antioxidant alpha-lipoic acid. *Free Rad. Biol. Med.* **22**, 359–378.

- Patlak C. S., Blasberg R. G. and Fenstermacher J. D. (1983) Graphical evaluation of blood-to-brain transfer constants from multiple-time uptake data. *J. Cereb. Blood Flow Metab.* **3**, 1–7.
- Pocernich C. B., Cardon A. L., Racine C. L., Lauderback C. M. and Butterfield D. A. (2002) Glutathione elevation and its protective role in acrolein-induced protein damage in synaptosomal membranes: relevance to brain lipid peroxidation in neurodegenerative disease. *Neurochem. Int.* **39**, 12–21.
- Preston E. and Hynie I. (1991) Transfer constants for blood–brain barrier permeation of the neuroexcitatory shellfish toxin, domoic acid. *Can. J. Neurol. Sci.* **18**, 39–44.
- Rapoport S. I. and Robinson P. J. (1995) Long-chain fatty acid transport at the blood–brain barrier and incorporation into brain phospholipids: a new *in vivo* method for examining neuroplasticity, and brain second messenger systems involving phospholipase A₂ activation, In: *New Concepts of a Blood–Brain Barrier* (Greenwood J., Begley D. J. and Segal M. B., eds), pp. 119–140. Plenum Press, New York.
- Rivas-Arancibia S., Dorado-Martinez C., Borgonio-Perez G., Hiriart-Urdanivia M., Verdugo-Diaz L., Duran-Vazquez A., Colin-Baranque L. and Avila-Costa M. R. (2000) Effects of taurine on ozone-induced memory deficits and lipid peroxidation levels in brains of young, mature, and old rats. *Environ. Res.* **82**, 7–17.
- Sato E., Kurokawa T., Oda N. and Ishibashi S. (1996) Early appearance of abnormality of microperoxisomal enzymes in the cerebral cortex of senescence-accelerated mouse. *Mech. Ageing Dev.* **92**, 175–184.
- Shih S. T., Khorram O., Lipton J. M. and McCann S. M. (1986) Central administration of α -MSH antiserum augments fever in the rabbit. *Am. J. Physiol.* **250**, R803–R806.
- Stadtman E. R. (1992) Protein oxidation and aging. *Science* **257**, 1220–1224.
- Stoll S., Hartmann H., Cohen S. A. and Muller W. E. (1993) The potent free radical scavenger α -lipoic acid improves memory in aged mice: putative relationship to NMDA receptor deficits. *Pharmacol. Biochem. Behav.* **46**, 799–805.
- Takeda T., Hosokawa M. and Higuchi K. (1991) Senescence-accelerated mouse (SAM): a novel murine model of accelerated senescence. *J. Am. Geriatr. Soc.* **39**, 911–919.
- Tanaka T., Yamada K., Senzaki K., Narimatsu H., Nishimura K., Kameyama T. and Nabeshima T. (1998) NC-1900, an active fragment analog of arginine vasopressin, improves learning and memory deficits induced by beta-amyloid protein in rats. *Eur. J. Pharmacol.* **352**, 135–142.
- Tappel A. L. (1968) Will antioxidant nutrients slow aging processes? *Geriatrics* **12**, 97–105.
- Umhauer S. A., Isbell D. T. and Butterfield D. A. (1992) Spin labeling of membrane proteins in mammalian brain synaptic plasma membranes: partial characterization. *Anal. Lett.* **25**, 1207–1215.
- Van Bree J. B. M. M., Tio S., De Boer A. G., Danhof M., Verhoef J. C. and Breimer D. D. (1990) Transport of desglycinamide-arginine vasopressin across the blood–brain barrier in rats as evaluated by the unit impulse response methodology. *Pharmacol. Res.* **7**, 293–298.
- Varadarajan S., Yatin S., Aksenova M. and Butterfield D. A. (2000) Review: Alzheimer's amyloid beta-peptide-associated free radical oxidative stress and neurotoxicity. *J. Structural Biol.* **130**, 184–208.
- Varadarajan S., Kanski J., Aksenova M., Lauderback C. and Butterfield D. A. (2001) Different mechanisms of oxidative stress and neurotoxicity for Alzheimer's A beta (1–43) and A beta (25–35). *J. Am. Chem. Soc.* **123**, 5625–5631.
- Wernerman J. and Hammarqvist F. (1996) Modulation of endogenous glutathione availability. *Curr. Opin. Clin. Nutri Metabolic Care* **2**, 487–492.
- Woods S. C. and Porte D. Jr (1983) The role of insulin as a satiety factor in the central nervous system, In: *Advances in Metabolic Disorders, CNS Regulation of Carbohydrate Metabolism*, Vol. 10 (Szabo A. J., ed.), pp. 457–482. Academic Press, New York.
- Yagi H., Katoh S., Akiguchi I. and Takeda T. (1988) Age-related deterioration of ability of acquisition in memory and learning in senescence accelerated mouse: SAM-P/8 as an animal model of disturbances in recent memory. *Neurosci. Biobehav. Rev.* **474**, 86–93.
- Yao C. Z., MacLellan D. G. and Thompson J. C. (1989) Intracerebroventricular administration of bombesin inhibits biliary and gastric secretion in the rat. *J. Neurosci. Res.* **22**, 461–463.
- Yatin S. M., Varadarajan S., Link C. D. and Butterfield D. A. (1999) In vitro and in vivo oxidative stress associated with Alzheimer's amyloid beta-peptide (1–43). *Neurobiol. Aging* **20**, 325–330.
- Yehuda S. and Youdim M. B. H. (1981) Effects of TRH and PS-24 on colonic temperature and motor activity of rats: possible role of dopamine. *Peptides* **2**, 131–135.
- Zs-Nagy I. (1978) A membrane hypothesis of aging. *J. Theor. Biol.* **75**, 189–195.
- Zs-Nagy I. (1990) Dietary antioxidants and brain aging: hopes and facts, In: *The Potential for Nutritional Modulation of Aging Processes* (Ingram D. K., Baker G. T. and Shock N. W., eds), pp. 379–399. Food and Nutrition Press, Trumbull.